

Stretched exponential relaxation in the mode-coupling theory for the Kardar-Parisi-Zhang equation

Francesca Colaioni and M. A. Moore

Department of Physics and Astronomy, University of Manchester, Manchester, M13 9PL, United Kingdom

(February 1, 2008)

We study the mode-coupling theory for the Kardar-Parisi-Zhang equation in the strong-coupling regime, focusing on the long time properties. By a saddle point analysis of the mode-coupling equations, we derive exact results for the correlation function in the long time limit – a limit which is hard to study using simulations. The correlation function at wavevector $\mathbf{k}$ in dimension d is found to behave asymptotically at time t as $C(\mathbf{k}, t) \simeq \frac{A}{k^{d+4-2z}} (Btk^z)^{\gamma/z} e^{-(Btk^z)^{1/z}}$, with $\gamma = (d-1)/2$, A a determined constant and B a scale factor.

PACS numbers: 05.40.-a, 64.60.Ht, 05.70.Ln, 68.35.Fx

The Kardar-Parisi-Zhang (KPZ) [1] equation is a simple non-linear Langevin equation, proposed in 1986 as a coarse grained description of a growing interface. Probably due to the fact that is the simplest generalization of the diffusion equation which includes a relevant non-linear term, the KPZ equation also arises in connection with many other important physical problems (the Burgers equation for one-dimensional turbulence [2], directed polymers in a random medium [3–5] etc.).

The KPZ equation for a growing interface, described by a single valued height function $h(\mathbf{x}, t)$ on a d -dimensional substrate $\mathbf{x} \in \mathbb{R}^d$ is:

$$\partial_t h(\mathbf{x}, t) = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta(\mathbf{x}, t). \quad (1)$$

The first term represents the surface tension forces which tend to smooth the interface, the second describes the nonlinear growth locally normal to the surface, and the last is a noise which mimics the stochastic nature of the growth process [6]. We choose the noise to be Gaussian, with zero mean and second moment $\langle \eta(\mathbf{x}, t) \eta(\mathbf{x}', t') \rangle = 2D \delta^d(\mathbf{x} - \mathbf{x}') \delta(t - t')$. The steady state interface profile is usually described in terms of the roughness: $w = \langle h^2(\mathbf{x}, t) \rangle - \langle h(\mathbf{x}, t) \rangle^2$ which for a system of size L behaves like $L^\chi f(t/L^z)$, where $f(x) \rightarrow \text{const}$ as $x \rightarrow \infty$ and $f(x) \sim x^{\chi/z}$ as $x \rightarrow 0$, so that w grows with time like $t^{\chi/z}$ until it saturate to L^χ when $t \sim L^z$. χ and z are the roughness and dynamic exponent respectively.

The phenomenology of the KPZ is well known: above two dimensions, there are two distinct regimes, separated by a critical value λ_c of the non-linearity coefficient. In the weak coupling regime ($\lambda < \lambda_c$) the non-linear term is irrelevant and the behavior is governed by the Gaussian ($\lambda = 0$) fixed point. The KPZ in this regime is equivalent to the linear Edward Wilkinson equation, for which the exponents are known exactly $\chi = (2-d)/2$ and $z = 2$. The more challenging strong coupling regime ($\lambda > \lambda_c$), where the non-linearity is relevant is characterised by anomalous exponents, which are not known exactly in general dimension d . From the Galilean in-

variance [2] (invariance of Eq. (1) under an infinitesimal tilting of the surface) one can derive the relation $\chi + z = 2$, which leaves just one independent exponent. For the special case when $d = 1$, the existence of a fluctuation-dissipation theorem gives the exact results $\chi = 1/2$, $z = 3/2$.

We take a self-consistent approach, the mode-coupling (MC) approximation [2], in which in the diagrammatic expansion for the correlation and response function only diagrams which do not renormalise the three point vertex λ are retained. The MC approximation has been remarkably successful in other areas of condensed matter physics, for example in the study of structural glass transitions [7], binary mixtures [8] and critical dynamics of magnets [8,9]. KPZ mode-coupling reproduces the exact values of the exponents in $d = 1$. Furthermore MC equations have been shown to arise from the large N -limit of an generalised N -component KPZ equation [10], which allows in principle a systematic approach to the theory beyond mode coupling, by expanding in $1/N$.

Our study focuses on the large time properties of the KPZ equation: we predict a stretched exponential decay for the correlation function at large times. A phenomenological stretched exponential law was used as long ago as 1854 to fit electronic relaxation data for a Leiden jar capacitor [11], and then been rediscovered many times: fitting functions involving stretched exponentials are nowadays widely used in phenomenological analysis of relaxation data (examples are dielectric relaxation [12] and glassy relaxation [7,13]). However, only a few analytical arguments [14] are able to reproduce stretched exponential relaxation in complex systems. Our prediction is in principle amenable of numerical verification, even though usual numerical techniques, mainly based on simulations [15] of discrete microscopical models which belong to the KPZ universality class [16], are hard in the long time asymptotic region.

Mode-coupling equations are coupled equation for the correlations and response function. The correlation and response function are defined in Fourier space by

$$C(\mathbf{k}, \omega) = \langle h(\mathbf{k}, \omega) h^*(\mathbf{k}, \omega) \rangle,$$

$$G(\mathbf{k}, \omega) = \delta^{-d}(\mathbf{k} - \mathbf{k}') \delta^{-1}(\omega - \omega') \langle \frac{\partial h(\mathbf{k}, \omega)}{\partial \eta(\mathbf{k}', \omega')} \rangle,$$

where $\langle \cdot \rangle$ indicate an average over η . In the mode-coupling approximation, the correlation and response functions are the solutions of two coupled equations,

$$G^{-1}(\mathbf{k}, \omega) = G_0^{-1}(\mathbf{k}, \omega) + \lambda^2 \int \frac{d\Omega}{2\pi} \int \frac{d^d q}{(2\pi)^d} [\mathbf{q} \cdot (\mathbf{k} - \mathbf{q})] [\mathbf{q} \cdot \mathbf{k}] G(\mathbf{k} - \mathbf{q}, \omega - \Omega) C(\mathbf{q}, \Omega), \quad (2)$$

$$C(\mathbf{k}, \omega) = C_0(\mathbf{k}, \omega) + \frac{\lambda^2}{2} |G(\mathbf{k}, \omega)|^2 \int \frac{d\Omega}{2\pi} \int \frac{d^d q}{(2\pi)^d} [\mathbf{q} \cdot (\mathbf{k} - \mathbf{q})]^2 C(\mathbf{k} - \mathbf{q}, \omega - \Omega) C(\mathbf{q}, \Omega), \quad (3)$$

where $G_0(\mathbf{k}, \omega) = (\nu k^2 - i\omega)^{-1}$ is the bare response function, and $C_0(\mathbf{k}, \omega) = 2D |G(\mathbf{k}, \omega)|^2$. In the strong coupling limit, $G(\mathbf{k}, \omega)$ and $C(\mathbf{k}, \omega)$ take the following scaling forms

$$G(\mathbf{k}, \omega) = k^{-z} g(\omega/k^z),$$

$$C(\mathbf{k}, \omega) = k^{-(2\chi+d+z)} n(\omega/k^z),$$

and Eqs. (2) and (3) translate into the following coupled equations for the scaling functions $n(x)$ and $g(x)$:

$$g^{-1}(x) = -ix + I_1(x), \quad (4)$$

$$n(x) = |g(x)|^2 I_2(x), \quad (5)$$

where $x = \omega/k^z$ and $I_1(x)$ and $I_2(x)$ are given by

$$I_1(x) = P \int_0^\pi d\theta \sin^{d-2} \theta \int_0^\infty dq \cos \theta (\cos \theta - q) q^{2z-3} r^{-z} \int_{-\infty}^\infty dy g\left(\frac{x - q^z y}{r^z}\right) n(y),$$

$$I_2(x) = \frac{P}{2} \int_0^\pi d\theta \sin^{d-2} \theta \int_0^\infty dq (\cos \theta - q)^2 q^{2z-3} r^{-(d+4-z)} \int_{-\infty}^\infty dy n\left(\frac{x - q^z y}{r^z}\right) n(y),$$

with $P = \lambda^2 / (2^d \Gamma(\frac{d-1}{2}) \pi^{(d+3)/2})$, $r^2 = 1 + q^2 - 2q \cos \theta$. It will be convenient to write Eqs. (4,5) as a function of time t , by Fourier transforming them:

$$\frac{\widehat{g_R}}{|g|^2}(t) = \widehat{I}_1(t) \quad (6)$$

$$\frac{\widehat{n}}{|g|^2}(t) = \widehat{I}_2(t) \quad (7)$$

where $\widehat{I}_1$ is the Fourier transform of the real part of I_1 and $\widehat{I}_2$ is the Fourier transform of I_2

$$\widehat{I}_1(t) = 2\pi P \int_0^\pi d\theta \sin^{d-2} \theta \int_0^\infty dq \cos \theta (\cos \theta - q) q^{2z-3} \widehat{g_R}(tr^z) \widehat{n}(tq^z),$$

$$\widehat{I}_2(t) = \pi P \int_0^\pi d\theta \sin^{d-2} \theta \int_0^\infty dq (\cos \theta - q)^2 q^{2z-3} r^{-(d+4-2z)} \widehat{n}(tr^z) \widehat{n}(tq^z).$$

Assuming an exponential, or stretched exponential decay for $\widehat{n}(t)$, we look for an asymptotic solution for $t \rightarrow \infty$ of Eq. (7) in the form

$$\widehat{n_\infty}(t) = A (Bt)^{\frac{\gamma}{2}} e^{-|Bt|^{\frac{\alpha}{2}}}. \quad (8)$$

We will show that such a solution exists and we will determine the exponents $\gamma = (d-1)/2$ and $\alpha = 1$. Even though the scaling part of the correlation and response function are coupled in the two Eqs. (6) and (7), we will see that the specific form of $\widehat{g}(t)$ is not relevant to the large t behaviour of $\widehat{n}(t)$. We will proceed as follows: we evaluate the right hand side of Eq. (7) in the large t limit by saddle point methods. The result of this analysis shows that any $\alpha \leq 1$ allows to reproduce the exponential factor. For $\alpha \leq z$, we can then assume that the left hand side of Eq. (7) is dominated by $g(0)^{-2} \widehat{n}(t)$ (see below), thus the asymptotic form of $I_2(t)$ has to be matched with $g(0)^{-2} \widehat{n_\infty}(t)$. The exponential factor can be matched by any $\alpha \leq 1$, but we will see that the only value of α that allows also to match the power law factor $t^{\gamma/z}$ is $\alpha = 1$. We can then proceed with $\alpha = 1$ to get $\gamma = \frac{d-1}{2z}$ and

$$A = \frac{g(0)^{-2} 4\Gamma(4z-4)}{P 2^{(d-1)/2} \Gamma((d-1)/2) \Gamma(2z-2)^2}. \quad (9)$$

Let us start with the evaluation of the left hand side of Eq. (7). Expanding $|g(x)|^2$ in even powers of x around $x = 0$, we can write $(n/\widehat{|g|}^2)(t) = a_0\widehat{n}(t) + a_1 d^2\widehat{n}(t)/dt^2 + \dots$ where $a_0 = g(0)^{-2}$. For $\widehat{n}(t)$ of the form (8) with $\alpha \leq z$, the first term of the series will dominate at large t , we can therefore assume that the left hand side of Eq. (7) has the asymptotic behaviour $(n/\widehat{|g|}^2)(t) \simeq g(0)^{-2}\widehat{n}(t)$. Let us now turn to the asymptotic analysis of $\widehat{I}_2(t)$. First note that the integral $\widehat{I}_2(t)$, is symmetric in the exchange $q \rightarrow r$, $\theta \rightarrow \phi$, where $r \sin \phi = q \sin \theta$ (see Fig 1). This allows us to rewrite $\widehat{I}_2(t)$ as twice the integral restricted to the region $q \cos \theta < 1/2$:

$$\widehat{I}_2(t) = 2\pi P \int_0^\infty dq \int_{\bar{\theta}}^\pi d\theta \sin^{d-2} \theta \int_0^\infty dq (\cos \theta - q)^2 q^{2z-3} r^{-(d+4-2z)} \widehat{n}(tr^z) \widehat{n}(tq^z), \quad (10)$$

where $\cos \bar{\theta} = \max[1/2q, 1]$. In $\widehat{I}_2(t)$ the function $\widehat{n}(t)$ appears in the integrand with the arguments tq^z , tr^z . For large enough t , we can always safely replace $\widehat{n}(tr^z)$ with its asymptotic form $\widehat{n}_\infty(tr^z)$, since in the new region of integration $r \geq 1/2$. We have to use more care with $\widehat{n}(tq^z)$, since such replacement is not allowed for all q , but only for $q > C/t^{1/z}$, where C is a large constant. We then have $\widehat{I}_2(t) \simeq \widehat{I}_2^\infty(t) = J_1(t) + J_2(t)$, where

$$J_1(t) = 2\pi P \int_0^{C/t^{1/z}} dq \int_{\bar{\theta}}^\pi d\theta \sin^{d-2} \theta (\cos \theta - q)^2 q^{2z-3} r^{-(d+4-2z)} \widehat{n}_\infty(tr^z) \widehat{n}(tq^z),$$

$$J_2(t) = 2\pi P \int_{C/t^{1/z}}^\infty dq \int_{\bar{\theta}}^\pi d\theta \sin^{d-2} \theta (\cos \theta - q)^2 q^{2z-3} r^{-(d+4-2z)} \widehat{n}_\infty(tr^z) \widehat{n}_\infty(tq^z).$$

The contribution from J_1 is negligible with respect to J_2 as $t \rightarrow \infty$. Let us evaluate an asymptotic expression for J_2 and postpone the discussion of J_1 to the end. Inserting $\widehat{n}_\infty(t) = A(Bt)^{\gamma/z} \exp(-|Bt|^{\alpha/z})$ in $J_2(t)$ gives

$$J_2(t) = 2\pi P A^2 (Bt)^{\frac{2\gamma}{z}} \int_{C/t^{1/z}}^\infty dq \int_{\bar{\theta}}^\pi d\theta \sin^{d-2} \theta (\cos \theta - q)^2 q^{2z-3+\gamma} r^{-(d+4-2z)+\gamma} e^{-(Bt)^{\frac{\alpha}{z}}(q^\alpha+r^\alpha)}. \quad (11)$$

It is immediately clear that the main contribution to this integral will come from the region where $f(q, \theta) = q^\alpha + r(q, \theta)^\alpha$ has its minimum, i.e. from the segment $\theta = 0$, $C/t^{1/z} < q < 1/2$ (for large enough t , $C/t^{1/z} < 1/2$). The value of q which minimises f will depend on α : for $\alpha < 1$, f reaches its minimum at $q_0 = C/t^{1/z}$, where $f(q_0, 0) = C^\alpha/t^{\alpha/z} + |1 - C/t^{1/z}|^\alpha \simeq 1 + (C/t^{1/z})^\alpha + O(C/t^{1/z})$. For $\alpha > 1$, the minimum is realized at $q = 1/2$, where $f(1/2, 0) = 1/2^{\alpha-1}$. For $\alpha = 1$, all q in the region $C/t^{1/z} < q < 1/2$ contribute equally, giving $f(q, 0) = 1$. The saddle point approximation in the angular integral gives:

$$J_2(t) \simeq 2\pi P A^2 (Bt)^{\frac{2\gamma}{z}} \int_{C/t^{1/z}}^{1/2} dq e^{-(Bt)^{\frac{\alpha}{z}}(q^\alpha+|1-q|^\alpha)} |1-q|^{2-(d+4-2z)+\gamma} q^{2z-3+\gamma} \int_0^\pi d\theta \theta^{d-2} e^{-(Bt)^{\frac{\alpha}{z}} \frac{\alpha}{2} q|1-q|^{\alpha-1} \theta^2}. \quad (12)$$

Performing the integral over θ in the large t limit:

$$J_2(t) \simeq \pi P A^2 \Gamma\left(\frac{d-1}{2}\right) \left(\frac{2}{\alpha}\right)^{\frac{d-1}{2}} (Bt)^{\frac{2\gamma}{z} - \frac{d-1}{2z}} \int_{C/t^{1/z}}^{1/2} dq e^{-(Bt)^{\frac{\alpha}{z}}(q^\alpha+|1-q|^\alpha)} |1-q|^\phi q^\psi, \quad (13)$$

where $\phi = 2z - d - 2 + \gamma + (1 - \alpha)\frac{d-1}{2}$ and $\psi = 2z - 3 + \gamma - \frac{d-1}{2}$. The leading contribution for large t is then given by

$$J_2(t) \simeq \pi P A^2 \Gamma\left(\frac{d-1}{2}\right) \left(\frac{2}{\alpha}\right)^{\frac{d-1}{2}} (Bt)^{\frac{2\gamma}{z} - \frac{d-1}{2z}} \begin{cases} \frac{\Gamma(\frac{\psi+1}{\alpha})}{\alpha} e^{-(Bt)^{\frac{\alpha}{z}}} (Bt)^{-\frac{\psi+1}{z}} & \alpha < 1 \\ I e^{-(Bt)^{\frac{1}{z}}} & \alpha = 1 \\ \sqrt{\pi} 2^{-(\phi+\psi)} e^{-\frac{(Bt)^{\frac{\alpha}{z}}}{2\alpha-1}} (Bt)^{-\frac{\alpha}{2z}} & \alpha > 1 \end{cases} \quad (14)$$

where $I = \int_0^{1/2} dq q^\psi (1-q)^\phi$. For $\alpha \leq 1$, the left hand side of Eq. (7) will be dominated by a term of the form $g(0)^{-2}\widehat{n}_\infty(t)$, thus $\widehat{I}_2(t) \simeq J_2(t)$ must be proportional to $\widehat{n}_\infty(t)$. This can only be achieved with $\alpha = 1$ and $\gamma = \frac{d-1}{2}$. For $\alpha < 1$, the stretched exponential factor is recovered, but trying to match the power of t leads to $\gamma = \frac{d-1}{2} + \psi + 1$, which can only be satisfied by the unphysical value $z = 1$. For $1 < \alpha < z$, the left hand side of Eq. (7) cannot be matched by J_2 . We then conclude that α must be equal to 1, and the asymptotic solution is then given by Eq. (8). The coefficient A can be determined as well by observing that $\gamma = \frac{d-1}{2}$ gives $\psi = \phi = 2z - 3$, and thus $I = \Gamma(2z-2)^2/2\Gamma(4z-4)$ which leads to Eq. (9).

Let us now go back to the integral J_1 . The argument of $\hat{n}(tq^z)$ in J_1 runs between 0 and C^z . Thus we can obtain an upper bound on J_1 by replacing $\hat{n}(tq^z)$ by its maximum in this region (which is given by $\hat{n}(0)$ if $\hat{n}(t)$ is a monotonically decreasing function). An analysis similarly to the one done for J_2 , can then be performed for J_1 , leading to $J_1 \propto (Bt)^{\frac{z}{z}-\frac{d-1}{2z}} e^{-(Bt)^{\frac{z}{z}}}$. Thus the contribution to $\hat{I}_2^\infty$ from J_1 is down by a factor $(Bt)^{-\frac{z}{z}}$.

The scale parameter B is just that, and remains unfixed in terms of the parameters of the KPZ equation. In a recent work [17], we proposed a scaling ansatz for the correlation function as $z \rightarrow 2$. If such an ansatz is correct, in a parametrization where $g(0)$ is finite the scale B^{-1} on which t varies would go to zero as z approaches 2: $B^{-1} \simeq (2 - z)$. Thus the asymptotic solution that we have found here could be read as an asymptotic solution for all t as z approaches 2.

In summary, we have presented an analytical derivation of an asymptotic solution of the mode-coupling equations for the strong coupling regime of the KPZ equation in the large t limit. We predict a stretched exponential relaxation for the correlation function, with a power law prefactor. We hope that this prediction will stimulate numerical investigations of long-time limits generally.

The authors acknowledge the support of EPSRC under grants GR/L38578 and GR/L97698.

-
- [1] M. Kardar, G. Parisi, and Y-C. Zhang, Phys. Rev. Lett. **56**, 889 (1986).
 - [2] D. Forster, D.R. Nelson, and M.J. Stephen Phys. Rev. A **16**, 732 (1977).
 - [3] D.A. Huse, C.L. Henley, and D.S. Fisher, Phys. Rev. Lett. **55**, 2924 (1985); M. Kardar and Y-C. Zhang Phys. Rev. Lett. **58**, 2087 (1987).

- [4] J.Z. Imbrie and T. Spencer, J. Stat. Phys. **52**, 609 (1988); J. Cook and B. Derrida, Europhys. Lett. **10**, 195 (1989); J. Cook and B. Derrida, J. Phys. A **23**, 1523 (1990); M.R. Evans and B. Derrida, J. Stat. Phys. **69**, 427 (1992).
- [5] G. Parisi J. Phys (Paris) **51**, 1595 (1990); M. Mezard J. Phys (Paris) **51**, 1831 (1990)
- [6] For a review see J. Krug and H. Spohn in *Solids far from equilibrium: Growth morphology and defects*, ed. by C. Godreche (Cambridge Univ. Press, Cambridge, 1992); A.-L. Barabasi and H. E. Stanley *Fractal Concepts in Surface Growth* (Cambridge Univ. Press, Cambridge, 1995); T. Halpin-Healy and Y-C. Zhang Phys. Rep. **254**, 215 (1995).
- [7] W. Götze, in *Liquids, Freezing, and the Glass Transition*, ed. J.P. Hansen, D. Levesque, and J. Zinn-Justin (Nort Holland, 1991). See also W. Götze and L. Sjögren, Rep. Prog. Phys. **55**, 241 (1992).
- [8] K. Kawasaki, Ann. Phys. (N. Y.) **61**, 1 (1970); K. Kawasaki, in *Phase Transitions and Critical Phenomena*, Vol 5a, ed. C. Domb and M. S. Green (Academic Press, New York, 1976).
- [9] E. Frey and F. Schwabl, Adv. Phys. **43**, 577 (1994).
- [10] J.P. Doherty, M.A. Moore, J.M. Kim, and A. Bray, Phys. Rev. Lett. **72**, 2041 (1994).
- [11] R. Kohlrausch, Poggendorf's Ann. Phys. **91**, 179 (1854).
- [12] G. Williams and D. C. Watts, Trans. Faraday Soc. **66**, 80 (1970).
- [13] V. Chamberlin, G. Mozurkewich, and R. Orbach, Phys. Rev. Lett. **52**, 867 (1984); F. Mezei and A. P. Murani, J. Magn. Mater. **14**, 211 (1979).
- [14] N. D. Donsker and S.R.S. Varadhan, Commun. Pure Appl. Math. **32**, 721 (1979); A. Bunde, S. Havlin, J. Klafter, G. Gräff, and A. Shechter, Phys. Rev. Lett. **78**, 3338 (1998); R.M.C. de Almeida, N. Lemke, and I.A. Campbell cond-mat/0003351
- [15] L. H. Tang Phys Rev. A **45**, 7162 (1992); T. Ala-Nissila, T. Hjelt, J.M. Kosterlitz, and O. Venäläinen, J. Stat. Phys. **72**, 207 (1993), E. Marinari, A. Pagnani and G. Parisi cond-mat/0005105.
- [16] M. Marsili, A. Maritan, F. Toigo, and J.R. Banavar Rev. Mod. Phys. **68**, 963 (1996).
- [17] F. Colaiori and M. A. Moore, cond-mat/0010410.




FIG. 1. Region of integration for $\hat{I}_2(t)$ in Eq. (10).